



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

Tnac
L72

JP



~~DEPOSITED AT THE~~
~~HARVARD FOREST~~
1943

RETURNED TO J. P.
MARCH, 1967



Dr George Little!

FOR THE PARIS EXPOSITION.

CATALOGUE
OF
ORES, ROCKS AND WOODS,

SELECTED FROM THE

Geological Survey Collection

OF THE

STATE OF GEORGIA, U. S. A.

WITH A DESCRIPTION OF THE GEOLOGICAL FORMATIONS.

GEORGE LITTLE, PH. D.,
STATE GEOLOGIST.

ATLANTA, GA.:

JAS. P. HARRISON & CO., STATE PRINTERS AND PUBLISHERS.

1878.



#

CATALOGUE
OF
ORES, ROCKS AND WOODS,

SELECTED FROM THE

Geological Survey Collection

OF THE

STATE OF GEORGIA, U. S. A.

WITH A DESCRIPTION OF THE GEOLOGICAL FORMATIONS.

GEORGE LITTLE, PH. D.,
STATE GEOLOGIST.

ATLANTA, GA.:

JAS. P. HARRISON & CO., STATE PRINTERS AND PUBLISHERS.

1878.

May 1903
18081

CATALOGUE.

The Geological Formations of Georgia embrace:

1. QUATERNARY, of the Atlantic Coast
Okefenokee Swamp.
Drift Deposits.
2. TERTIARY, Eocene of the..... Vicksburg group.
" Jackson group.
" Claiborne group.
3. SECONDARY, Cretaceous Nos. 5, 4 and 1 of.. Upper Missouri section.
4. PALEOZOIC, of the Carboniferous Period.... Coal-measures.
" Millstone grit.
" Sub-carb. limestone.
Devonian..... Black Shale—Genesee.
" Oriskany.
Upper Silurian..... Niagara.
" Clinton.
" Medina.
Lower Silurian..... Cincinnati.
" Trenton.
" Chazy.
" Quebec.
" Potsdam.
5. METAMORPHIC.—No fossils. Stratigraphic equivalents of Lower Silurian. Subdivisions not all distinguished.

ROCKS AND MATERIAL, REPRESENTING FORMATIONS.

QUATERNARY AGE.

White and yellow sands, with drift (water-worn pebbles); white and variegated clays; argillaceous sandstones (white clay and sand conglomerate); coarse white micaceous sands; yellowish sands, with ferruginous sandstones, ferr. concretions and concretionary iron ore.

TERTIARY PERIOD.

Red and yellow loam, with concretionary iron ore—brown hematite; limestone; buhrstone; greenish white clays, with loose silicified shells; greensand (glauconite); white joint clays, with lignitic seams and ledges of claystone; white clay marls; silicious fossiliferous sandstone; interstratified sands and shaly clay; white sandy marl; compact clayey limestone (indurate marl); blue fossiliferous joint-clay; blue and gray shell marls, with lignite; compact white and yellow limestones; white

friable coral limestone; white joint clay; hard and compact white limestone (in some cases arenaceous).

CRETACEOUS PERIOD.

Limestone, highly fossiliferous; blue shell-marl, alternating with nodular limestone strata; blue micaceous glauconitic clay; yellow fossiliferous clay; blue micaceous clay, with few fossils; white clays and sands.

CARBONIFEROUS AGE.

Coal-measures.—Sandstones, shales, ironstones, fireclays, and three to five beds of coal. 500 feet.

Millstone grit.—Conglomerates, sandstones, shales, ironstones, and one bed of coal. 400 feet.

Upper Sub-carboniferous.—Massive limestones, partly oölitic, mostly pure, less lime to the eastward, few shaly beds. 500 feet.

Lower Sub-carboniferous.—Silicious group of Safford; limestone, more or less impure, mostly filled with silicious concretions; at some points shales, partly bituminous and thin-bedded argillaceous limestones. 200 feet.

DEVONIAN AGE.

Genesee Epoch.—Black shale, bituminous; pyritous sandstone at base. 100 feet.

UPPER SILURIAN AGE.

Oriskany Epoch.—Very silicious limestone, weathering to a skeleton on the outcrop. 4 feet.

Clinton Epoch.—Shales, few thin limestones and argillaceous sandstones, and one to three beds of fossiliferous hematite. 400 feet.

Medina Epoch.—Ferruginous sandstones, few shales. 400 feet.

LOWER SILURIAN AGE.

Cincinnati Epoch.—Thin-bedded, ferruginous, silicious limestones and shales; hematites, rarely of workable thickness. 200 to 400 feet.

Trenton Epoch.—Thin-bedded limestones and shales, changing eastward to ferruginous shales. 700 feet.

Chazy Epoch.—Limestone, mostly shaly, becoming more ferruginous and argillaceous to the eastward. 600 feet.

Quebec Epoch.—Upper part—Cherty magnesian limestone, generally heavy-bedded, with one or two layers of white saccharoidal sandstone near middle of limestone. 400 feet. Lower part—Shales, often calcareous, with few generally thin-bedded limestones, partly oölitic. 2,000 feet.

Calceiferous Epoch.—Shales, often glauconitic, thin-bedded sandstones, rarely glauconitic, and silicious argillaceous limestones. 2,000 feet.

Potsdam Epoch.—Hard, ferruginous sandstones, generally quartzitic, partly thick-bedded, partly thin and argillaceous. 2,000 (?) feet.

Acadian Epoch.—Upper part—Everywhere partly metamorphosed slates, with conglomerate layers; passing into schists and gneisses. 10,000 (?) feet. Lower part—Hydromica schists. 2,000 feet.

METAMORPHIC.

Apparently all are stratigraphical equivalents of the Lower Silurian; but the subdivisions are not all distinguished. No fossils found.

Cincinnati Group.—Gneisses, largely hornblendic, sometimes quartzitic.

Few hydromica schists. Often includes epidiosytes (?) [see No. 190]. Some copper, little gold. Probably includes the flexible sandstones of the upper part of the "Itacolumyte Series" of Lieber. 15,000 (?) feet thick.

Quebec Group.—At top, limestones and dolomytes, with chert more or less altered into tremolite, accompanied by heavy beds of chrysolite, serpentine, chlorite and steatyte, often with abundant crystals and masses of corundum, chromite, spinel, etc. Some gold, galenite and traces of silver. The lower part of the group consists largely of hydromica schists, with thick and thin beds of gneiss, hornblende schists, quartzytes and partly solidified sandstones. Includes the lower part of Lieber's "Itacolumyte Series." Gold indigenous, abundant; less copper; much garnet, cyanite, rutile, ilmenite, magnetite, pyrite; a few diamonds. 12,000 (?) feet thick.

Upper Potsdam Group (Calciferous not separately recognized)—Heavy-bedded gneiss, with few schists; little hornblende. Thin steatytes, with serpentine and asbestus, at top. Little or no gold, often pyritous. 2,000 (?) feet thick.

Lower Potsdam, or Acadian.—Mica and hydromica schists; little hornblende. Bands of gneiss, often conglomeratic. Some gold, workable copper; much pyrite; some roofing slate. 13,000 (?) feet thick.

LIST OF SPECIMENS EXHIBITED.

ORES AND PRODUCTS.

1. Lead Ore—Galenite.....	Lincoln county.	
2. Manganese Ore—Pyrolusite.....	Lincoln	..
3. Copper Ore—Chalcopyrite, etc.....	Paulding	..
4. Copper Ore.....	Paulding	..
5. Sulphur Ore—Pyrite.....	Fulton	..
6. Sulphur Ore—Pyrite	Paulding	..
7. Sulphur Ore—Pyrite	Haralson	..
8. Chrome Ore.	Troup	..
9. Iron Ore—Hematite.....	Carroll	..
10. Iron Ore—Hematite	Bartow	..
11. Iron Ore—Limonite.....	Gordon	..
12. Iron Ore—Limonite.....	Bartow	..
13. Iron.....	Rome Iron Co., Floyd	..
14. Steel Rail	Atlanta Rolling Mill, Fulton	..
15. Spiegel Iron	Pool's Furnace, Bartow	..
16. Iron.....	Atlanta Rolling Mill, Fulton	..
17. Ferro-manganese.....	Ward's Furnace, Bartow	..
18. Manganese Ore—Pyrolusite	Bartow	..
18a. Manganese Ore.....	Bartow	..
19. Asbestos.....	Fulton	..
19a. Asbestos.....	Rabun	..
20. Heavy Spar—Barite.....	Bartow	..
21. Corundum.....	Carroll	..
22. Corundum in Serpentine.....	Towns	..
22A. Gold Ore.....	Lumpkin	..
22B. Gold Ore.....	Lumpkin	..
22C. Gold Ore	Lumpkin	..
22D. Gold Ore.....	Lumpkin	..
22E. Gold Ore	White	..
22F. Gold Ore.....	Paulding	..

BUILDING STONES, ETC.

23. Sandstone	Bartow	..
24. Limestone.....	Bartow	..
24a. Lime (made from No. 24).....	Bartow	..
25. White Limestone.....	Polk	..
26. Marble.....	Haralson	..
27. Marble.....	Whitfield	..
28. Marble, fossiliferous.	Walker	..
29. Black Marble.....	Polk	..
30. Black Marble.....	Whitfield	..
31. White Marble (3 specimens).....	Pickens	..
32. Variegated Marble	Pickens	..
33. Fire Brick	Baldwin	..

GEOLOGICAL SERIES.

NAME.	FORMATION.	GROUP.	COUNTY.
42. Opal.....	Tertiary.....	Jackson.....	Washington.
43. Agate.....	Tertiary.....	Jackson.....	Washington.
45. Buhrstone.....	Tertiary.....	Jackson.....	Calhoun.
46. Sandstone, fossiliferous.....	Tertiary.....	Jackson.....	Jefferson.
47. Sandstone, fossiliferous.....	Tertiary.....	Jackson.....	Wilkinson.
48. Silicified Wood.....	Quaternary.....	Clay.
49. Sandstone, ferruginous.....	Quaternary.....	Houston.
50. Sandstone, fossiliferous.....	Tertiary.....	Jackson.....	Jefferson.
51. Sandstone, ferr. argillaceous.....	Quaternary.....	Dodge.
52. Sandstone, fossiliferous.....	Tertiary.....	Jackson.....	Glascok.
53. Iron Ore.....	Tertiary.....	Jackson.....	Stewart.
54. Clay, white.....	Tertiary.....	Jackson.....	Washington.
55. Limestone.....	Tertiary.....	Claiborne.....	Clay.
56. Limestone.....	Tertiary.....	Passage beds—Jackson and Claiborne.....	Houston.
57. Marl, yellow.....	Tertiary.....	Jackson.....	Burke.
58. Marl, white.....	Tertiary.....	Screven.
59. Marl, yellow.....	Tertiary.....	Claiborne.....	Clay.
60. Greensand marl.....	Tertiary.....	Jackson.....	Houston.
61. Calcite.....	Tertiary.....	Jackson.....	Twiggs.
62. Limestone.....	Tertiary.....	Vicksburg.....	Mitchell.
63. Flint, ferruginous.....	Tertiary.....	Pulaski.
64. Buhrstone.....	Tertiary.....	Burke.
65. Limestone.....	Tertiary.....	Vicksburg(?).....	Pulaski.
66. Gypsum.....	Cretaceous.....	Stewart.
67. Shell marl, blue.....	Cretaceous.....	Clay.
68. Coal.....	Carboniferous.....	Dade.
68a. Coal.....	Carboniferous.....	Walker.
69. Sandstone.....	Carboniferous.....	Dade.
70. Sandstone.....	Carboniferous.....	Dade.
71. Conglomerate.....	Carboniferous.....	Walker.
72. Concretion.....	Sub-carboniferous.....	Walker.
73. Concretion.....	Sub-carboniferous.....	Floyd.
74. Concretion.....	Sub-carboniferous.....	Catoosa.
75. Limestone, chert partly dec'd.....	Sub-carboniferous.....	Walker.
76. Limestone.....	Sub-carboniferous.....	Gordon.
77. Chert, oolitic.....	Sub-carboniferous.....	Walker.
78. Chert.....	Sub-carboniferous.....	Walker.
79. Shale, blue, with concretions.....	Devonian.....	Genesee.....	Catoosa.
80. Shale, black.....	Devonian.....	Genesee.....	Walker.
81. Chert.....	Upper Silurian.....	Oriskany.....	Catoosa.
82. Hematite, fossiliferous.....	Upper Silurian.....	Clinton.....	Dade.
83. Hematite, fossiliferous.....	Upper Silurian.....	Clinton.....	Dade.
84. Hematite, fossiliferous.....	Upper Silurian.....	Clinton.....	Dade.
85. Hematite, fossiliferous.....	Upper Silurian.....	Clinton.....	Chattooga.
86. Hematite, fossiliferous.....	Upper Silurian.....	Clinton.....	Walker.
87. Sandstone, argillaceous.....	Upper Silurian.....	Clinton.....	Dade.
88. Sandstone, ferruginous.....	Upper Silurian.....	Medina.....	Whitfield.
89. Conglomerate, fine.....	Upper Silurian.....	Medina.....	Whitfield.
90. Sandstone, ferruginous.....	Lower Silurian.....	Cincinnati.....	Whitfield.
91. Sandstone.....	Lower Silurian.....	Cincinnati.....	Whitfield.
92. Limestone.....	Lower Silurian.....	Cincinnati.....	Whitfield.
93. Sandstone, ferruginous.....	Lower Silurian.....	Cincinnati.....	Walker.
94. Limestone.....	Lower Silurian.....	Trenton.....	Chattooga.
95. Limestone.....	Lower Silurian.....	Trenton.....	Walker.
96. Concretion, silicious.....	Lower Silurian.....	Chazy.....	Walker.
97. Sandstone, polishing.....	Lower Silurian.....	Upper Quebec.....	Whitfield.
98. Sandstone, saccharoidal.....	Lower Silurian.....	Upper Quebec.....	Whitfield.
99. Breccia.....	Lower Silurian.....	Upper Quebec.....	Catoosa.
100. Chert.....	Lower Silurian.....	Upper Quebec.....	Catoosa.
101. Chert, cellular.....	Lower Silurian.....	Upper Quebec.....	Walker.
102. Chert, concretion.....	Lower Silurian.....	Upper Quebec.....	Walker.
103. Chert, oolitic.....	Lower Silurian.....	Upper Quebec.....	Gordon.
104. Chert, with quartz veins.....	Lower Silurian.....	Upper Quebec.....	Gordon.
105. Chert.....	Lower Silurian.....	Upper Quebec.....	Walker.
106. Limestone, cherty.....	Lower Silurian.....	Upper Quebec.....	Walker.
107. Limestone, variegated.....	Lower Silurian.....	Upper Quebec.....	Whitfield.
108. Limestone.....	Lower Silurian.....	Upper Quebec.....	Whitfield.
109. Limestone, red variegated.....	Lower Silurian.....	Upper Quebec.....	Walker.
110. Limestone, dolomitic.....	Lower Silurian.....	Upper Quebec.....	Catoosa.
111. Limestone, oolitic.....	Lower Silurian.....	Lower Quebec.....	Catoosa.

GEOLOGICAL SERIES.—Continued.

NAME.	FORMATION.	GROUP.	COUNTY.
112. Shale, glauconitic.....	Lower Silurian.....	Lower Quebec.....	Whitfield.
113. Limestone, with galenite	Lower Silurian.....	Calceiferous.....	Catoosa.
114. Schist, chloritic.....	Lower Silurian.....	Calceiferous.....	Catoosa.
115. Limestone, silicious.....	Lower Silurian.....	Calceiferous.....	Chattooga.
116. Quartzite.....	Lower Silurian.....	Potsdam	Bartow.
117. Quartzite.....	Lower Silurian.....	Potsdam.....	Bartow.
118. Gneiss.....	Lower Silurian.....	Acadian.....	Murray.
119. Gneiss, porphyritic.....	Lower Silurian.....	Acadian.....	Bartow.
120. Conglomerate.....	Lower Silurian.....	Acadian.....	Bartow.
121. Conglomerate.....	Lower Silurian.....	Acadian.....	Bartow.
122. Conglomerate.....	Lower Silurian.....	Acadian.....	Gilmer.

METAMORPHIC ROCKS.

123. Flint.....	Lower Silurian	Quebec	Lincoln.
124. Quartzite.....	Lower Silurian	Cincinnati.....	Towns.
125. Quartzite.....	Lower Silurian	Cincinnati.....	Hall.
126. Quartzite, pyritous.....	Harris.
127. Quartzite, pyritous.....	Lower Silurian.....	Quebec.....	White.
128. Schist, silicious, auriferous.....	Lower Silurian.....	Quebec.....	White.
129. Schist, silicious, auriferous.....	Lower Silurian.....	Quebec.....	White.
130. Quartzite, schistose, auriferous	Lower Silurian	Quebec.....	White.
131. Itacolumyte.....	Lower Silurian	Cincinnati (?).....	Hall.
132. Itacolumyte.....	Lower Silurian	Cincinnati (?).....	Hall.
133. Itacolumyte.....	Lower Silurian	Quebec (?).....	Heard.
134. Gneiss, quartzitic, garnetiferous	Lower Silurian	Cincinnati (?).....	Talbot.
135. Quartzite, cellular	Lower Silurian	Quebec.....	Hall.
136. Quartzite, gneissoid	Lower Silurian	Potsdam	Heard.
137. Quartzite, gneissoid	Lower Silurian	Potsdam	Muscogee.
138. Granyte, pink.....	Lower Silurian.....	Muscogee.
139. Feldspar	Lower Silurian.....	Cincinnati.....	Habersham.
140. Kaolin	Lower Silurian.....	Cincinnati (?).....	Union.
141. Kaolin	Lower Silurian.....	Cincinnati (?).....	Fulton.
142. Quartzite, conglomeratic.....	Lower Silurian.....	Cincinnati.....	Gilmer.
143. Quartzite, conglomeratic.....	Lower Silurian.....	Acadian.....	Bartow.
144. Conglomerate, granytoid.....	Lower Silurian.....	Acadian.....	Bartow.
145. Gneiss, quartzitic.....	Lower Silurian.....	Cincinnati.....	Lumpkin.
146. Granite.....	Lower Silurian.....	Cincinnati (?).....	DeKalb.
147. Gneiss.....	Lower Silurian.....	Cincinnati.....	Fannin.
148. Gneiss, banded.....	Lower Silurian.....	Quebec.....	Fulton.
149. Gneiss.....	Lower Silurian.....	Cincinnati.....	Habersham.
150. Gneiss, biotitic, porphyritic.....	Lower Silurian.....	Acadian (?).....	Harris.
151. Gneiss, biotitic.....	Lower Silurian.....	Cincinnati (?).....	Campbell.
152. Gneiss, porphyritic.....	Lower Silurian.....	Cincinnati (?).....	Habersham.
153. Gneiss, biotitic, porphyritic.....	Lower Silurian.....	Quebec	Lumpkin.
154. Schist silicious, ferruginous, decayed	Lower Silurian.....	Quebec.....	White.
155. Mica schist.....	Lower Silurian.....	Quebec.....	Hart.
156. Hydromica schist.....	Lower Silurian.....	Cincinnati (?).....	Towns.
157. Hydromica schist, decayed.....	Lower Silurian.....	Calceiferous (?).....	Gilmer.
158. Mica schist.....	Lower Silurian.....	Cincinnati.....	Habersham.
159. Argillite, decayed.....	Lower Silurian.....	Quebec.....	Gilmer.
160. Argillite, silicious.....	Lower Silurian.....	Cincinnati.....	Gilmer.
161. Hydromica schist porphyritic.....	Lower Silurian.....	Cincinnati.....	Gilmer.
162. Hydromica schist, garnetiferous	Lower Silurian.....	Calceiferous.....	Gilmer.
163. Hydromica schist, garnetiferous	Lower Silurian.....	Quebec.....	Lumpkin.
164. Hydromica schist, garnetiferous	Lower Silurian.....	Quebec.....	Lumpkin.
165. Chlorite schist, graphitic.....	Lower Silurian.....	Cincinnati.....	Gilmer.
166. Chlorite schist.....	Lower Silurian.....	Quebec.....	Habersham.
167. Chlorite schist.....	Lower Silurian.....	Quebec.....	Habersham.
168. Chlorite schist.....	Lower Silurian.....	Quebec.....	Douglas.
169. Steatyte.....	Lower Silurian.....	Quebec.....	DeKalb.
170. Steatyte.....	Lower Silurian.....	Quebec.....	Rabun.
171. Talcose schist, silicious.....	Lower Silurian.....	Quebec.....	Taylor.
172. Hornblende slate, talcoid	Lower Silurian.....	Quebec.....	Lumpkin.

METAMORPHIC ROCKS.—*Continued.*

NAME.	FORMATION	GROUP.	COUNTY.
173. Hornblende schist	Lower Silurian....	Quebec.....	Lumpkin.
174. Hornblende schist, decayed..	Lower Silurian....	Quebec.. ..	Lumpkin.
175. Hornblende schist, decayed..	Lower Silurian....	Quebec.....	White.
176. Hornblende schist, pyritous..	Lower Silurian....	Quebec.....	Towns.
177. Hornblende schist, pyritous..	Lower Silurian....	Quebec.....	Towns.
178. Hornblendic gneiss	Lower Silurian....	Cincinnati.....	Towns.
179. Hornblendic gneiss	Lower Silurian....	Cincinnati.....	Rabun.
180. Hornblendic gneiss, decayed..	Lower Silurian....	Cincinnati (?).....	Towns.
181. Syenitic gneiss	Lower Silurian....	Cincinnati.....	Lumpkin.
182. Mica schist, calcareous	Lower Silurian....	Cincinnati.....	Gilmer.
183. Limestone, shaly, pyritous..	Lower Silurian....	Cincinnati.....	Gilmer.
184. Limestone, micaceous.....	Lower Silurian....	Cincinnati.....	Gilmer.
185. Chert	Lower Silurian....	Quebec.....	Hall.
186. Tremolite, calcareous, (al- tered chert)	Lower Silurian....	Quebec.....	Hall.
187. Quartzite, cyanitic.....	Lower Silurian ..	Quebec.....	Lumpkin.
188. Quartzite, chrysolitic.....	Lower Silurian....	Quebec.....	Dawson.
189. Dunyte.....	Lower Silurian....	Quebec.....	White.
190. ? Epidosite ?.....	Lower Silurian....	Cincinnati.....	Towns.
191. Trap.....	Jurassic (?).....	Coweta.
192. Trap.....	Jurassic (?).....	Habersham.
193. Trap.....	Jurassic (?).....	DeKalb.
194. Trap.....	Jurassic (?).....	DeKalb.
195. Trap, decayed.	Jurassic (?).....	Franklin.
196. Trap, decayed.....	Jurassic (?).....	Gwinnett.
197. Trap.....	Jurassic (?).....	Lincoln.

WOODS.

198. *Magnolia glauca*—Sweet Bay.....Okefenokee swamp.
199. *Liriodendron tulipifera*—Poplar.....Murray county.
200. *Æsculus pavia*—Buckeye Whitfield ..
201. *Acer saccharinum*—Sugar Maple..... Whitfield ..
202. *Acer dasycarpum*—Silver Maple..... Murray ..
203. *Prunus serotina*—Wild Cherry..... Murray ..
204. *Prunus spinosa*—Sloe..... Walker ..
205. *Liquidambar styraciflua*—Sweet Gum..... Murray ..
206. *Nyssa multiflora*—Sour Gum. Murray ..
207. *Ilex opaca*—Holly..... Murray ..
208. *Diospyros Virginiana*—Persimmon..... Murray ..
209. *Persea Carolinensis*—Red Bay..... Okefenokee swamp.
210. *Sassafras officinalis*—Sassafras Murray county.
211. *Ulmus Americana*—Elm..... Murray ..
212. *Ulmus fulva*—Slippery Elm..... Murray ..
213. *Ulmus alata*—Wahoo Murray ..
214. *Celtis occidentalis*—Hack-berry..... Walker ..
215. *Platanus occidentalis*—Sycamore..... Whitfield ..
216. *Carya tomentosa*—Hickory..... Whitfield ..
217. *Carya alba*—Shell-bark Hickory..... Whitfield ..
218. *Juglans nigra*—Black Walnut..... Murray ..
219. *Castanea Americana*—Chestnut..... Whitfield ..
220. *Fagus ferruginea*—Beech. Murray ..
221. *Ostrya Virginica*—Hop Hornbean..... Murray ..
222. *Salix nigra*—Black Willow..... Whitfield ..
223. *Pinus glabra*—Spruce Pine..... Murray ..
224. *Pinus rigida*—Pitch Pine..... Murray ..
225. *Juniperus Virginiana*—Red Cedar..... Walker ..
226. *Taxodium distichum*—Cypress Okefenokee swamp.
227. *Sabal palmetto*—Cabbage Palmetto..... Glynn county.

GEOLOGICAL DESCRIPTION.

The general strikes of the Georgia outcrops being approximately northeast and southwest, a section from northwest to southeast shows most readily the general structure.

The extreme northwest corner of the State includes a small area of the Coal-measures, nearly level and undisturbed, forming, practically, an outlier of the great plateau of the Cumberland Mountains. But a small part of the series exposed in these mountains appears here, and this probably belongs to the Millstone grit. Starting southeastward, we almost immediately encounter, along the edges of the plateau outliers, upturned portions of the same beds, in which the first signs of metamorphism appear, in the partially de-bitumenized coals.

The Subcarboniferous beds have mostly very limited outcrops, as mere borders of the small Coal-measure areas, except along one line of outliers from which the Coal-measures have been entirely removed, and only Subcarboniferous occupies the surface.

The upper limestones are mostly quite pure, and yield fossils in some abundance. They seem to represent the Chester and St. Louis divisions of the formation. The lower limestones are very impure, containing large amounts of cherty concretions, with comparatively few fossils. The shales at the base of the series are partly bituminous, partly silicious, and contain, at bottom, the band of phosphatic nodules which mark the horizon through Tennessee, Kentucky and Southern Indiana.

The Devonian black shale, of the Genesee epoch, varies from ten to eighty feet, or more, in thickness, and is of no economical importance. Few fossils have been found, either in the shale itself, or in the pyritous, shaly sandstone at its base, whose age is still in doubt, since it may be properly part of the shale group, or it may represent some other portion of the Hamilton group, just as a pyritous fish-bearing shale represents the Tully limestone of New York, in the western counties of that State.

Of the Oriskany, but a few inches have been identified by fossils, and that only at a single locality. This shows a lenticular mass of quartzose sandstone, from which the calcareous fossils have been weathered out, along the outcrop.

The Clinton group presents several distinct lines of outcrop, of a thick series of variously colored, mostly ferruginous, calcareous shales, with a few thin beds of limestone and argillaceous sandstone, which are sometimes crowded with fossils. In its western outcrops, it includes three distinct beds of oölitic hematite, more or less calcareous, all frequently of workable thickness; but to the eastward these are reduced to a single layer, and that finally but a few inches thick. The sandy layers increase in this direction.

The thick ferruginous sandstone of the Medina outcrops along ridge lines, making marked features of the landscape. Passing the last of

these, we finally leave behind us all the newer Paleozoic rocks, and enter a region of exclusively Lower Silurian strata, stretching to the southern and eastern edges of the metamorphic area. These lower beds have also appeared in the several anticlinal valleys already crossed, but mostly not in so full development as where we now find them.

The Cincinnati group consists mainly of thin bedded limestones, which are pretty pure to the westward, but become increasingly silicious and ferruginous, and thickens somewhat as we pass eastward. In a few localities, they are quite shaly. At many points, there are small deposits of fossiliferous oolitic hematite, rarely, however, in sufficient amount to be workable. In the more easterly outcrops, these ores are frequently more or less metamorphosed into "specular" ore, sometimes distinctly crystallized. The ferruginous, silicious limestones have also there suffered incipient metamorphism—the structure lines being much distorted, the lime carbonate partially crystallized, and the whole mass solidified to almost a quartzitic texture, while the fossils are still distinct, and the iron is still mechanically, rather than chemically, combined.

The Trenton group is represented by a thin series of limestones, pure to the westward, becoming argillaceous and ferruginous, and finally shales, to the eastward. The shales are hardly to be distinguished from those above and below, within the non-metamorphic area, and become utterly undistinguishable in their metamorphic condition.

The Chazy limestones are more or less shaly through their entire outcrop; but the impurities of the western portion become the main constituent of the layers in their eastern extension.

The Quebec group dolomites are heavy-bedded, and of enormous thickness—commonly reaching a total of from 3,000 to 4,000 feet. With the exception of occasional layers, they are very impure throughout their entire outcrop, being commonly argillaceous and silicious, and often inclosing nodules and masses of chert in immense quantity. They very rarely yield fossils. Neither in degree of impurity nor in thickness, do they vary much, in passing from one part of the area to another. The shales of the lower part of the group are generally calcareous, and include a few thin beds of fossiliferous limestone. They vary but little in composition, but are partially altered into slates in their eastern portions.

Of the area thus far described, the western portion rises into sharp ridges, reaching elevations of from 1,200 to 2,300 feet above the sea, with intervening valleys at levels of from 700 to 1,000 feet. The eastern portion, occupying the flatter half of the continuation of the "Great Valley" of East Tennessee and Virginia, has an average elevation of about 1,000 feet for the ridges, and perhaps 800 for the valleys. It may be noted here, that the higher ridges are all synclinals, while the anticlinals are marked by the lowest valleys. In leaving this non-metamorphic area, bounded by the Cohutta mountains on the east, and

by the Dugdown mountain on the south, we rise sharply, except along the gap occupied by the Western and Atlantic Railroad, from 800 to about 3,000 feet above the level of the adjoining portions of the "Great Valley." The actual limit of the area we are leaving, is a line of faults, on the northwestern side of which only a slight degree of metamorphism is found; while, on the opposite side, nearly everything is much altered. The first rocks encountered are commonly the semi-metamorphic slates, assigned, with little doubt, to the Acadian, and provisionally termed the "Ocoee" by Safford.

They occupy long but generally narrow areas. Immediately above them, we commonly find the heavy-bedded ferruginous sandstones of the Potsdam (Chilhowee of Safford,) not seen within the non-metamorphic area, which is mostly quartzitic along the Cohutta range, but is represented further eastward and southward by gneisses of all grades of crystallization. Along only one line of outcrop do we find any evidence of rocks underlying and older than the Acadian. This single exception is the area of soft hydromica schists, with quartz veins, which includes the copper region running south from the southeast corner of Tennessee. As these strata show no signs of unconformability, we are obliged to account them part of the Acadian group.

In these outer portions of the metamorphic area, the alteration of the beds is but partial, the shales being baked into argillites, more or less glassy; the sandstones solidified into quartzites, with the sand grains and pebbles still distinct; the limestones crystallized, but with their impurities only partially combined. But as we pass southeastward, the change becomes more complete, the shales becoming mica, hydromica, chlorite and talc schists; the chert masses of the Quebec group are crystallized into tremolite; the magnesian shales and limestones become steatites, serpentines and chrysolytes (dunyte); the sandstones become quartzites and gneisses, often losing all lamination, and appearing as granites; the quartz veins, which are at first few in number, become numerous, and are partially replaced by veins of granite. These latter are at first small, but are finally very large, and even appear to be the sole occupants of extensive areas. Here again the synclinals commonly form the mountains and higher ridges, while the lines of anticlinals are commonly marked by valleys.

The entire mass of the metamorphic rocks, so far as yet definitely determined, is of Lower Silurian age. Much the larger part of the area has been identified, beyond a question, with the Potsdam, Quebec and Cincinnati groups, leaving only very limited patches within which it is still possible that small islands of Archean rocks may be recognized. As already stated, the Trenton, Chazy and Calcareous beds vary so slightly from the Quebec and Potsdam strata that they can be distinguished with difficulty, even when not metamorphosed; and not at all when altered, at least with the amount of study which it has yet been possible to bestow upon them. We are, therefore, compelled to neglect them as being but subordinate fractions of the larger groups. The Quebec

group, already 6,000 feet thick in the non-metamorphic area, now presents a possible maximum of from 12,000 to 15,000 feet, and forms much the larger portion of the metamorphic area. It is also, economically, the most important portion, because it is the matrix of nearly all the gold of Georgia. It also includes workable deposits of copper ore, and large amounts of corundum. A few diamonds have been found upon its outcrops. The Cincinnati group occupies, generally, narrow areas, and yields but little mineral wealth, including small amounts of gold, copper and iron. The Potsdam outcrops are of still smaller extent, and yield almost nothing except building stone. The Acadian sub-group yields roofing slate and copper ores.

Crossing the metamorphic rocks, generally along lines approximately transverse to the strike, there are several dykes of various forms of trap. Their formation evidently occurred after the upturning and metamorphism of the enclosing strata, which disturbance is referred to the Permian; and they are probably of the same age as the Jurassic (?) dykes which accompany the Triassic red sandstones of North Carolina and more northern States.

CRETACEOUS.

This formation has but a small representation in Georgia. Its beds are exposed in a narrow belt extending from Columbus southward, nearly to Fort Gaines, and consist of blue micaceous marls and greensand clays, yellow fossiliferous with white plastic clays.

An abundance of fossils, both vertebrate and invertebrate, exist in the blue and yellow marls, and the beds are highly pyritiferous, with limestone nodules and gypsum. The country is usually broken from denudation, and the hills capped with sand and ferruginous sandstone.

The marls contain about ten per cent. of lime, with potash and phosphoric acid, averaging, each, about .5 per cent. The greensand clays have about six per cent. of pure greensand, the thickness of the bed being from fifteen to twenty feet, and are exposed in the river bluffs.

TERTIARY.

The tertiary marls of the State are confined mainly to a belt of country, averaging about forty miles in width, and lying immediately south of a line drawn across the State from Fort Gaines, through Macon and Milledgeville, to Augusta.

The lowest and thickest bed is a white friable marl, containing, at some localities, as high as ninety per cent. of carbonate of lime. In the other beds, the proportion of carbonate of lime varies from this to a mere trace. Along this Eocene outcrop, there are also deposits of almost pure "green sand," or glauconite, while we find this mineral, at various localities, present also in the calcareous marl, in almost every possible proportion. These marls have been used to considerable extent as fertilizers; and in almost every instance the result has been highly satisfactory. Where it has been otherwise, the fault has been either in the quantity used, or in the mode of application.

The buhrstone also extends entirely across the State, and overlies the calcareous beds. Some very fine mill-stones have been manufactured from this rock, in Burke and Jefferson counties.

In the Tertiary loams—as well as in the superficial quaternary deposits—there occurs a good quality of concretionary iron ore—brown hematite. This has been worked in some localities.

QUATERNARY.

This formation is found capping the hills and overlying the Tertiary beds mentioned above, and may be considered as co-extensive therewith. Its useful materials are—

1. **CLAYS.**—These are sedimentary in character, and vary from pure white porcelain-clay, to dark purplish pipe-clay. They are already being utilized, in the manufacture of crockery ware, pottery, fire-brick, etc.; and, from their excellent quality, their utilization is destined to become one of the leading industries of the State.

2. **SANDSTONES, OR CLAY AND SAND CONGLOMERATES.**—These are also sedimentary—the finer material above them having been longer held in suspension, and deposited more quietly. They are composed of white clay, and coarse sand or gravel. They harden on exposure, and make a first-class building material.

3. **FERRUGINOUS SANDSTONES.**—The ferruginous sandstones are very abundant, and of good quality. They are easily quarried, and make a most excellent building stone.

TIMBER.

Nearly all that portion of the State lying south of the line drawn above, as the northern boundary of the Tertiary, is timbered with long-leaf *yellow pine*. This section embraces about thirty thousand square miles. Supposing twenty per cent. of this to be cleared, (which is an over-estimate,) we have twenty-four thousand square miles of pine timber, which will produce, on an average, 30,000 feet of first-class *heart* lumber *per acre*.

WATER POWERS OF GEORGIA.

The principal water powers of Georgia are found north of a line passing through Columbus, Macon and Augusta. The waters of the branches, creeks and small rivers of this elevated region are accumulated several hundred feet above the ocean level, and precipitated from the tough metamorphic rocks upon the lower Tertiary country below, thus forming some of our most gigantic water powers, at an elevation entirely free from malaria, and immediately surrounded by the cotton growing regions of Carolina, Georgia and Alabama.

The estimates given below are for the theoretical horse-power of the stream, without the accumulation of its waters in a reservoir. The horse-power is equivalent to 33,000 foot-pounds.

Chattahoochee river, Columbus.....	35,552 horse-power.
Chattahoochee river, Fulton county	2,448 horse-power.
Ocmulgee river, Lloyd's shoals.....	8,970 horse-power.
Ocmulgee river, Seven Islands ...	2,040 horse-power.
Ocmulgee river, Capps Shoals.....	508 horse-power.
Ocmulgee river, Glover's mill	1,368 horse-power.
Etowah river, Bartow county.....	2,250 horse-power.
Etowah river, Franklin mines.....	1,029 horse-power.
Etowah river, Lumpkin county	272 horse-power.
Holt's Shoal's Bibb county.....	1,050 horse-power.
South River, Butts county.	850 horse-power.
South River, Clark's Factory.....	247 horse-power.
Snake creek, Carroll county.....	405 horse-power.
Pataula creek, Clay county.....	601 horse-power.
Armuchee creek, Floyd county.....	151 horse-power.
Coosawattee, Carter's mill.....	3,085 horse-power.
Oconee river, Long Shoals Factory.....	1,824 horse-power.
Oconee river, Riley's shoals	2,054 horse-power.
Oconee river, Oconee county.....	5,642 horse-power.
Oconee river, Jackson county.....	271 horse-power.
Tallulah river, Habersham county.....	20,508 horse-power.
Mulberry creek, Harris county.....	1,020 horse-power.
Towaliga, High Falls.....	1,530 horse-power.
Yellow River, Cedar Shoals.....	1,302 horse-power.
Yellow River, Cedar and Henley shoals.....	2,000 horse-power.
Little River, Eatonton Factory.....	155 horse-power.
Nacoochee Gold Mining Co., White county.....	575 horse-power.
Savannah river, Augusta canal.....	12,000 horse-power.

These are only a few of the many which might be mentioned.



